

Journal Pre-proof

Increases in 'deaths of despair' during the COVID-19 pandemic in the USA and UK

Colin Angus, MSc, Mr, Charlotte Buckley, PhD, Dr, Andrea M. Tilstra, PhD, Dr,
Jennifer B Dowd, PhD, Professor



PII: S0033-3506(23)00079-3

DOI: <https://doi.org/10.1016/j.puhe.2023.02.019>

Reference: PUHE 4837

To appear in: *Public Health*

Received Date: 20 January 2023

Revised Date: 20 February 2023

Accepted Date: 22 February 2023

Please cite this article as: Angus C, Buckley C, Tilstra AM, Dowd JB, Increases in 'deaths of despair' during the COVID-19 pandemic in the USA and UK, *Public Health*, <https://doi.org/10.1016/j.puhe.2023.02.019>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2023 The Author(s). Published by Elsevier Ltd on behalf of The Royal Society for Public Health.

Increases in 'deaths of despair' during the COVID-19 pandemic in the USA and UK

Mr Colin Angus^{*1,2} MSc, Dr Charlotte Buckley³ PhD, Dr Andrea M. Tilstra² PhD, Professor Jennifer B. Dowd² PhD

¹School of Health and Related Research, University of Sheffield, UK

²Leverhulme Centre of Demographic Science, Department of Sociology, University of Oxford, UK

³Department of Automatic Controls and Systems Engineering, University of Sheffield, UK

*Corresponding author. Email c.r.angus@sheffield.ac.uk

Declaration of competing interest: Nothing to declare

Funding: We acknowledge funding from the Leverhulme Trust Large Centre Grant, Leverhulme Centre for Demographic Science (AMT, JBD) and the European Research Council grant ERC-2021-CoG-101002587 (AMT, JBD), the John Fell Fund, University of Oxford (CA, JBD) and the National Institute on Alcohol Abuse and Alcoholism of the National Institutes of Health under Award Number R01AA028009 (CB). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

Abstract

Objectives

The COVID-19 pandemic significantly impacted mental health, health-related behaviours such as drinking and illicit drug use, and the accessibility of health and social care services. How these pandemic shocks affected “despair”-related mortality in different countries is less clear. This study uses public data to compare deaths from alcohol, drugs and suicide in the USA and UK to identify similarities or differences in the impact of the pandemic on important non-COVID causes of death across countries and to consider the public health implications of these trends.

Study design and methods

Data was taken from publicly available mortality figures for England & Wales, Northern Ireland, Scotland and the United States of America 2001-2021 and analysed descriptively through age-standardised and age-specific mortality rates from suicide, alcohol and drug use.

Results

Alcohol-specific deaths increased in all countries between 2019 and 2021, most notably in the USA and, to a lesser extent, England & Wales. Suicide rates did not increase markedly during the pandemic in any of the included nations. Drug-related mortality rates rose dramatically over the same period in the USA but not in other nations.

Conclusions

Mortality from ‘deaths of despair’ during the pandemic has displayed divergent trends between causes and countries. Concerns about increases in deaths by suicide appear to have been unfounded, while deaths due to alcohol have risen across the UK and in the USA and across almost all age groups. Scotland and the USA had similarly high levels of drug-related deaths pre-pandemic, but the differing trends during the pandemic highlight the different underlying causes of these drug death epidemics and the importance of tailoring policy responses to these specific contexts.

Keywords

COVID-19, Public Health, Alcohol, Drugs, Suicide, Mortality

Introduction

The global COVID-19 pandemic dramatically changed daily life for most people in high-income countries. The early months of the pandemic saw the closure of schools, bars and restaurants, and severe restrictions on social interactions. The resulting social isolation raised concerns about potential increases in depression and suicide.¹ Similar concerns were voiced about increases in heavy drinking and substance misuse to cope with stress of the pandemic. Subsequent data showed significant deteriorations in mental health during the pandemic, particularly among women, young people and those on low incomes.² Prior to the pandemic, increasing levels of mortality from alcohol, drugs and suicide – so called ‘deaths of despair’ – had been well documented in the US.³ Some have identified this as a uniquely American phenomenon,⁴ however disaggregating the constituent nations of the UK highlights some dramatic trends. In recent years Scotland has seen drug-related deaths and male suicide rates rise on a par with increases seen in the US, while alcohol-related deaths have fallen. At the same time deaths attributable to alcohol have risen consistently in England and Northern Ireland.⁵

While tax data suggested that there was no notable change in overall alcohol consumption in the UK during the first two years of pandemic,⁶ individual-level surveys showed an increase in heavier drinking, suggesting greater polarisation of drinking behaviour in England,^{7,8} although not necessarily in Scotland.⁹ Survey data in the US showed more drinking days per month and heavier drinking, particularly for women.¹⁰ The pandemic also led to significant changes in accessibility of services including mental health care and specialist alcohol and drug treatment services that have benefitted some groups, but restricted access to others.^{11,12}

Whether the dire predictions around increased mortality from alcohol, drugs and suicide – so-called ‘deaths of despair’ played out during the pandemic is not clear. Some studies suggest little impact on suicide rates,¹³ but notable increases in alcohol deaths in the UK¹⁴ in 2020 as well as substantial increases in alcohol and drug-related mortality in the US.^{15,16} We use publicly available mortality data from 2001 and 2021 to compare mortality rates from these three causes across nations to better understand the wider impacts of the pandemic on public health.

Methods

We used mortality data for England & Wales from the ‘21st Century Mortality File’ published by the Office for National Statistics,¹⁷ for Northern Ireland from the Annual Reports of the Registrar General published by the Northern Ireland Statistics and Research Agency,¹⁸ for Scotland from the Vital Events Reference Tables published by National Records of Scotland¹⁹ and for the USA from the Centers for Disease Control and Prevention’s WONDER Underlying Cause of Death data (for 2001-2020) and Multiple Cause of Death data (provisional) (for 2021).²⁰ Data was available by ICD-10 code, sex and in 5-year age bands for all countries. Deaths were categorised based on the underlying cause of death as being attributable to alcohol, drugs or suicide in non-overlapping categories on the basis of standard ICD-10 code definitions (see supplementary table S1 for the list of included ICD-10 codes).^{5,21} Age-standardised mortality rates per 100,000 population for each cause from 2000-2021 were calculated using the European Standard Population²² and population estimates from the Human Mortality Database.²³

Results

Age-standardised mortality rates for each cause, by sex and country are illustrated in Figure 1 and summarised in Table 1. In all countries, suicide rates did not change much during the pandemic period. In the USA, rates for both men and women dipped slightly *below* the pre-pandemic trend, while there was a small increase for both men and women in Northern Ireland. Suicide rates in England & Wales and Scotland were essentially stable. Changes in alcohol-specific deaths during the pandemic were more noticeable. Mortality increased for both men and women in England & Wales and the USA. In England & Wales, women saw an increase from 10.4 to 12.9 per 100,000 in alcohol-specific deaths from 2019 to 2021, a relative increase of 24.1%, compared to an increase of 19.6% in men. The USA saw even greater increases in alcohol-specific deaths of 29.1% in women and 26.7% in men, and rates rose to a lesser extent for both men and women in Northern Ireland. Deaths from alcohol increased among Scottish men (23.7% relative increase) after a consistently declining pre-pandemic trend. The picture for drug-related deaths was more mixed, with a continuation of the gradually rising pre-pandemic trend in England & Wales and Northern Ireland. Both Scotland and the USA had seen sharp rises in drug-related deaths in the years immediately prior to the pandemic, but this trend levelled off in Scotland, while accelerating dramatically in the USA, particularly among men. Drug-related deaths among men in the USA rose from 33.5 to 51.6 per 100,000 from 2019 to 2021, a 54.1% relative increase. For women in the USA the increase was also substantial, from 14.4 to 21.2 per 100,000 (47.9% increase).

Figure 1 and Table 1 here

We examined differences in these trends by age in Figure 2. In England & Wales alcohol-specific deaths rose for everyone except the youngest age groups, while increases in drug-related deaths were largely confined to 45-64 year-old age groups that had seen rising pre-pandemic rates of drug-related mortality. There are few clear age patterns for Northern Ireland, although there is some evidence of an increase in alcohol deaths among those over age 65. In Scotland, alcohol-specific deaths rose sharply in men over 45, reversing sharp declines for many years prior to the pandemic. While drug-related mortality in Scotland had risen steeply in recent years, deaths fell for those aged 35-44 during the pandemic but continued to rise among 45-54 year-old men. Finally, age-specific data for the USA paints an alarming picture, with both alcohol and drug deaths rising sharply across almost every age group.

Figure 2 here**Discussion**

The COVID-19 pandemic disrupted daily life on an unprecedented scale, closing schools and businesses and restricting social interactions for many months. Concerns about the mental health impacts of social isolation and other disruptions were raised early on, but it's unclear how much this translated into increased non-COVID mortality. We summarised mortality from three "despair"-related causes (alcohol, drug-related, suicide) in the USA and UK nations over the pandemic and compared them to pre-pandemic trends. Despite these concerns, we found little apparent association between the pandemic on deaths by suicide. The most consistent increase across countries was in alcohol-specific deaths. The USA stood alone in its dramatic increase in drug-related deaths, compounding already very high levels. England & Wales and Northern Ireland saw slight increases in drug-related deaths in line with upward previous trends, while changes in drug deaths in Scotland were flat despite a strong upward trend in recent years. These patterns were largely consistent across age groups, with no evidence that younger age groups suffered more.

The COVID-19 pandemic was expected to impact substance use for many reasons. COVID disrupted most aspects of daily life and increased social isolation, potentially increasing demand for alcohol and drugs to cope with stress. Supply was also impacted by the closures of typical venues for social drinking such as pubs and restaurants. A priori, the closing of spaces for social drinking could reduce overall drinking if drinking happens less often at home. On the other hand, the pandemic was a shock to people's routines, and working from home could have made it easier to drink at home. Evidence from England and Wales does suggest shifts towards more heavy drinking during the pandemic,^{7,8,24} but this is less clear in Scotland.⁹ In the USA and UK sales of alcohol spiked in March 2020 in anticipation of stay-at-home orders,^{10,25} and surveys suggested more drinking and heavy drinking days in the months that followed. People may have substituted alcohol for legal or illegal drugs that were harder to obtain during the pandemic.²⁶ The consistent increase in alcohol-specific deaths across countries suggests that levels of heavy drinking did increase during the pandemic. Most alcohol-specific deaths are due to alcohol-related liver disease,²⁷ which typically develops over many years,²⁸ in contrast to poisoning which is more acute but makes up a small percentage of alcohol-specific deaths. Given the sharp increase in alcohol deaths since the pandemic, this suggests that the pandemic induced extra drinking among already heavy drinkers who were near the threshold of succumbing to liver disease.

It is also likely that access to treatment and harm reduction services was reduced during the time of pandemic restrictions, increasing the potential lethality of these behaviours.²⁹ Drug supply shortages

can also drive consumption of riskier substances and behaviours such as sharing injecting equipment.³⁰ People may also purchase larger quantities of drugs at one time when they have the opportunity, increasing the risk of overdose. The extent and nature of these issues will vary substantially depending on the local context and the types of drugs implicated in drug-related deaths. In particular we might expect them to differ between the US, where rising drug-related deaths are primary linked to prescription opioids and, more recently, fentanyl,³¹ and Scotland where the rise in drug-related deaths is strongly linked to 'street' benzodiazepines.³²

There are some limitations to our analyses. ICD-10 code-based definitions are imperfect, and use of codes may vary between countries. Our definitions of each cause of death are in line with previous, similar, studies,²¹ but these do not align completely with existing definitions of 'alcohol-specific/induced' deaths as used in the UK and USA. We exclude several, generally rare, causes such as alcoholic polyneuropathy (ICD-10 code G62.1) as these cannot be disaggregated in the public mortality data we have used, while we have included hepatitis and cirrhosis of the liver (ICD-10 codes K73-74), of which approximately 70% of deaths are estimated to be due to alcohol.³³ A comparison of these two definitions shows close alignment of trends (see supplementary material). This approach remains likely to significantly underestimate the true burden of alcohol mortality as we exclude other conditions for which alcohol is a contributing cause, such as cancers and cardiovascular disease.³⁴ For drug-related and suicide mortality our approach yields similar figures to the official 'drug poisoning'/'drug-induced causes' and suicide figures published within each country (see supplementary material), with differences accounted for largely by the fact that we have defined our causes of death to be non-overlapping, whereas deaths from deliberate overdoses, for example, may be included in published figures for both suicide and drug-related deaths. Our approach does not account for deaths where multiple causes are implicated, for example combined alcohol and opioid poisonings, which have been shown to account for a substantial minority of all deaths from alcohol and drug poisoning in the US.³⁵ Finally, it should be noted that the relatively smaller populations of Northern Ireland and, to a lesser extent Scotland, mean that data for these countries is inherently subject to greater random variation year-on-year and greater caution should therefore be exercised when interpreting annual fluctuations.

Our analysis of age-standardised mortality trends provides straightforward description of changes in these rates during the pandemic. We do not estimate excess mortality by cause, which could more formally incorporate previous trends but is beyond the scope of this analysis. The USA observed substantial differences in drug-related mortality by racial and ethnic group over the pandemic, with overdose mortality particularly accelerating in Black relative to White groups.³⁶ Although beyond the scope of this study, this would be important to incorporate in future explorations of drug-related mortality. Finally, we did not consider socioeconomic differences in 'deaths of despair,' although it has been widely documented that these deaths are clustered in those with lower socioeconomic status, particularly in the United States,³ and it is likely that the pandemic widened existing socioeconomic gaps in mortality.³⁷ Further investigation of these socioeconomic and ethnic differences in mortality trends during the pandemic, and how these may be linked to aspects of the COVID response and wider government policy may help to understand the full impacts of the pandemic on health inequalities.

The data presented here paints a mixed picture of the impact of the pandemic on 'deaths of despair' in the USA and UK nations. There is little evidence of an increase in deaths by suicide, a marked increase in deaths due to alcohol and notably different trends in drug-related deaths between countries. The increases in alcohol and drug misuse deaths in the USA are particularly stark and suggest that policy action is urgently required. However, the many-faceted interactions between

pre-pandemic trends, the existing policy landscape, the direct impact of the pandemic and pandemic response measures, and the broader societal consequences of these mean that it is vital any public health policy response is tailored to the specific circumstances of both the country and affected populations.

The fact that we have seen further increases in mortality rates in 2021 on top of sharp increases in 2020 for alcohol and, in some cases, drug misuse deaths may suggest that these trends are linked more strongly to wider societal impacts of the pandemic, rather than the short-term policy responses to the initial COVID wave in March 2020. Whether these increases in mortality will return to pre-pandemic levels in the coming years represents a major public health concern, and understanding their underlying drivers is an important challenge that may inform short-term policy responses and guide planning for future pandemics.

Data availability and sharing

All data analysed in this study is publicly available. Code to process this data and reproduce the analysis presented here can be found at <https://github.com/VictimOfMaths/DeathsOfDespair/blob/master/DataInsight/DodPandemicPaperFinal.r>

Declaration of interests

Nothing to declare

References

1. Akinin LB, Andretti B, Goldszmidt R, Helliwell JF, Petherick A, Neve JED, et al. Policy stringency and mental health during the COVID-19 pandemic: a longitudinal analysis of data from 15 countries. *The Lancet Public Health*. 2022 May 1;7(5):e417–26.
2. McGinty EE, Presskreischer R, Han H, Barry CL. Trends in Psychological Distress Among US Adults During Different Phases of the COVID-19 Pandemic. *JAMA Network Open*. 2022 Jan 24;5(1):e2144776.
3. Case A, Deaton A. Rising morbidity and mortality in midlife among white non-Hispanic Americans in the 21st century. *Proceedings of the National Academy of Sciences*. 2015 Dec 8;112(49):15078–83.
4. Sterling P, Platt ML. Why Deaths of Despair Are Increasing in the US and Not Other Industrial Nations—Insights From Neuroscience and Anthropology. *JAMA Psychiatry*. 2022 Apr 1;79(4):368–74.
5. Dowd JB, Angus C, Zajacova A, Tilstra AM. Midlife “Deaths of Despair” Trends in the US, Canada, and UK, 2001-2019: Is the US an Anomaly? [Internet]. medRxiv; 2022 [cited 2022 Nov 11]. Available from: <https://www.medrxiv.org/content/10.1101/2022.10.10.22280916v1>
6. HM Revenue & Customs. Alcohol Bulletin [Internet]. GOV.UK. [cited 2022 Nov 11]. Available from: <https://www.gov.uk/government/statistics/alcohol-bulletin>
7. Jackson SE, Garnett C, Shahab L, Oldham M, Brown J. Association of the COVID-19 lockdown with smoking, drinking and attempts to quit in England: an analysis of 2019–20 data. *Addiction*. 2021;116(5):1233–44.

8. Jackson SE, Beard E, Angus C, Field M, Brown J. Moderators of changes in smoking, drinking and quitting behaviour associated with the first COVID-19 lockdown in England. *Addiction* [Internet]. 2021;117(3). Available from: <http://onlinelibrary.wiley.com/doi/abs/10.1111/add.15656>
9. Stevely AK, Sasso A, Hernandez Alava M, Holmes J. Changes in alcohol consumption in Scotland during the early stages of the COVID-19 pandemic: Descriptive analysis of repeat cross-sectional survey data [Internet]. Edinburgh: Public Health Scotland; 2021 p. 86. Available from: <https://www.publichealthscotland.scot/media/2983/changes-in-alcohol-consumption-in-scotland-during-the-early-stages-of-the-covid-19-pandemic.pdf>
10. Pollard MS, Tucker JS, Green HD Jr. Changes in Adult Alcohol Use and Consequences During the COVID-19 Pandemic in the US. *JAMA Network Open*. 2020 Sep 29;3(9):e2022942.
11. Schofield J, Dumbrell J, Matheson C, Parkes T, Bancroft A. The impact of COVID-19 on access to harm reduction, substance use treatment and recovery services in Scotland: a qualitative study. *BMC Public Health*. 2022 Mar 15;22(1):500.
12. Cordis Bright. Flexible responses during the Coronavirus crisis: rapid evidence gathering [Internet]. Make Every Adult Matter; 2020 [cited 2022 Dec 15]. Available from: <http://meam.org.uk/wp-content/uploads/2020/06/MEAM-Covid-REG-report.pdf>
13. Pirkis J, John A, Shin S, DelPozo-Banos M, Arya V, Analuisa-Aguilar P, et al. Suicide trends in the early months of the COVID-19 pandemic: an interrupted time-series analysis of preliminary data from 21 countries. *The Lancet Psychiatry*. 2021 Jul 1;8(7):579–88.
14. Holmes J, Angus C. Alcohol deaths rise sharply in England and Wales. *BMJ*. 2021 Mar 5;372:n607.
15. Hedegaard H, Miniño A, Spencer MR, Warner M. Drug Overdose Deaths in the United States, 1999–2020 [Internet]. National Center for Health Statistics (U.S.); 2021 Dec [cited 2022 Dec 13]. Available from: <https://stacks.cdc.gov/view/cdc/112340>
16. White AM, Castle IJP, Powell PA, Hingson RW, Koob GF. Alcohol-Related Deaths During the COVID-19 Pandemic. *JAMA*. 2022 May 3;327(17):1704–6.
17. Office for National Statistics. Deaths registered in England and Wales – 21st century mortality [Internet]. [cited 2022 Nov 11]. Available from: <https://www.ons.gov.uk/peoplepopulationandcommunity/birthsdeathsandmarriages/deaths/datasets/the21stcenturymortalityfilesdeathsdataset>
18. Northern Ireland Statistics and Research Agency. Registrar General Annual Report [Internet]. Northern Ireland Statistics and Research Agency. [cited 2022 Nov 11]. Available from: <https://www.nisra.gov.uk/statistics/births-deaths-and-marriages/registrar-general-annual-report>
19. National Records of Scotland. Vital Events Reference Tables [Internet]. National Records of Scotland. National Records of Scotland; [cited 2022 Nov 11]. Available from: <https://www.nrscotland.gov.uk/statistics-and-data/statistics/statistics-by-theme/vital-events/general-publications/vital-events-reference-tables>
20. Centre for Disease Control and Prevention. CDC WONDER [Internet]. [cited 2022 Nov 11]. Available from: <https://wonder.cdc.gov/Deaths-by-Underlying-Cause.html>


21. Masters RK, Tilstra AM, Simon DH. Explaining recent mortality trends among younger and middle-aged White Americans. *International Journal of Epidemiology*. 2018 Feb 1;47(1):81–8.
22. Pace M, Lanzieri G, Glickman M, Grande E, Zupanic T, Wojtyniak B, et al. Revision of the European Standard Population - Report of Eurostat's task force - 2013 edition [Internet]. Eurostat; [cited 2022 Nov 11]. Available from: <https://data.europa.eu/doi/10.2785/11470>
23. Max Planck Institute for Demographic Research (Germany), University of California, Berkeley (USA), and French Institute for Demographic Studies (France). HMD. Human Mortality Database [Internet]. Available from: www.mortality.org
24. Angus C, Henney M, Pryce R. Modelling the impact of changes in alcohol consumption during the COVID-19 pandemic on future alcohol-related harm in England [Internet]. The University of Sheffield; 2022 Jul [cited 2022 Nov 11]. Available from: https://figshare.shef.ac.uk/articles/report/Modelling_the_impact_of_changes_in_alcohol_consumption_during_the_COVID-19_pandemic_on_future_alcohol-related_harm_in_England/19597249/1
25. Finlay I, Gilmore I. Covid-19 and alcohol—a dangerous cocktail. *BMJ*. 2020 May 20;369:m1987.
26. United Nations Office on Drugs and Crime. COVID-19 and the drug supply chain: from production and trafficking to use [Internet]. Vienna; 2020 [cited 2022 Dec 14]. Available from: <https://www.unodc.org/documents/data-and-analysis/covid/Covid-19-and-drug-supply-chain-Mai2020.pdf>
27. Office for National Statistics. Alcohol-specific deaths in the UK 2021 [Internet]. 2022 [cited 2022 Dec 14]. Available from: <https://www.ons.gov.uk/peoplepopulationandcommunity/birthsdeathsandmarriages/deaths/bulletins/alcohol-specific-deaths-in-the-uk/2021-registrations>
28. Osna NA, Donohue TM, Kharbanda KK. Alcoholic Liver Disease: Pathogenesis and Current Management. *Alcohol Res*. 2017;38(2):147–61.
29. Jacka BP, Janssen T, Garner BR, Yermash J, Yap KR, Ball EL, et al. Impacts of the COVID-19 pandemic on healthcare access among patients receiving medication for opioid use disorder. *Drug Alcohol Depend*. 2021 Apr 1;221:108617.
30. Otiashvili D, Mgebrishvili T, Beselia A, Vardanashvili I, Dumchev K, Kiriazova T, et al. The impact of the COVID-19 pandemic on illicit drug supply, drug-related behaviour of people who use drugs and provision of drug related services in Georgia: results of a mixed methods prospective cohort study. *Harm Reduction Journal*. 2022 Mar 9;19(1):25.
31. Ciccarone D. The triple wave epidemic: Supply and demand drivers of the US opioid overdose crisis. *International Journal of Drug Policy*. 2019 Sep 1;71:183–8.
32. McAuley A, Matheson C, Robertson J. From the clinic to the street: the changing role of benzodiazepines in the Scottish overdose epidemic. *International Journal of Drug Policy*. 2022 Feb 1;100:103512.
33. Jaccard, A, Blawat A, Ashton S, Belsman L, Gommon J, Webber L, et al. Alcohol-attributable fractions for England: An update [Internet]. London: Public Health England; 2020. Available from:

https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/958648/RELATI_1-1.pdf



34. Rehm J, Gmel Sr GE, Gmel G, Hasan OSM, Imtiaz S, Popova S, et al. The relationship between different dimensions of alcohol use and the burden of disease—an update. *Addiction*. 2017;112(6):968–1001.
35. Buckley C, Ye Y, Kerr WC, Mulia N, Puka K, Rehm J, et al. Trends in mortality from alcohol, opioid, and combined alcohol and opioid poisonings by sex, educational attainment, and race and ethnicity for the United States 2000–2019. *BMC Med*. 2022 Oct 24;20(1):405.
36. Friedman JR, Hansen H. Evaluation of Increases in Drug Overdose Mortality Rates in the US by Race and Ethnicity Before and During the COVID-19 Pandemic. *JAMA Psychiatry*. 2022 Apr 1;79(4):379–81.
37. Stokes AC, Lundberg DJ, Elo IT, Hempstead K, Bor J, Preston SH. COVID-19 and excess mortality in the United States: A county-level analysis. *PLOS Medicine*. 2021 May 20;18(5):e1003571.

Table 1: Changes in age-standardised rates of 'deaths of despair' mortality 2019-2021.



Country	Sex	Cause	Age-standardised deaths per 100,000			Change 2019-21	
			2019	2020	2021	Absolute	Relative
England & Wales	Male	Alcohol	20.1	22.8	24.0	3.9	19.6%
		Drugs	10.0	10.5	11.0	1.0	9.9%
		Suicide	14.2	13.5	14.2	0.0	0.1%
	Female	Alcohol	10.4	12.1	12.9	2.5	24.1%
		Drugs	4.3	4.3	4.6	0.3	7.8%
		Suicide	4.1	4.0	4.7	0.6	14.0%
Northern Ireland	Male	Alcohol	27.0	31.1	29.6	2.7	9.9%
		Drugs	13.8	15.8	15.8	2.0	14.3%
		Suicide	17.9	19.1	20.0	2.1	11.7%
	Female	Alcohol	16.1	17.5	17.4	1.4	8.4%
		Drugs	4.8	5.8	5.5	0.7	15.0%
		Suicide	5.1	6.9	6.8	1.7	33.8%
Scotland	Male	Alcohol	30.0	36.0	37.1	7.1	23.7%
		Drugs	38.8	41.5	39.8	1.0	2.5%
		Suicide	21.9	20.9	22.2	0.2	1.0%
	Female	Alcohol	16.3	15.7	18.1	1.8	10.9%
		Drugs	16.3	14.9	15.9	-0.4	-2.6%
		Suicide	6.8	7.5	6.8	0.0	-0.1%
USA	Male	Alcohol	29.4	34.7	37.3	7.9	26.7%
		Drugs	33.5	44.4	51.6	18.1	54.1%
		Suicide	26.9	26.0	27.1	0.2	0.7%
	Female	Alcohol	13.6	16.2	17.6	4.0	29.1%
		Drugs	14.4	18.3	21.2	6.9	47.9%
		Suicide	6.7	6.1	6.3	-0.3	-5.2%





England & Wales





Northern Ireland



Scotland



USA



Journal Pre-proof