
# Colombia


Total population (2016): 48 654 392 > Population aged 15 years and older (15+): 76% > Population in urban areas: 77% > Income group (World Bank): Upper-middle income

## ALCOHOL CONSUMPTION: LEVELS AND PATTERNS

Recorded alcohol per capita (15+) consumption, 1961–2016



Recorded alcohol per capita (15+) consumption (in litres of pure alcohol) by type of alcoholic beverage, 2016 or latest year available



Alcohol per capita (15+) consumption (in litres of pure alcohol)

	20	10*	20	16*
Recorded	4.3		4.4	
Unrecorded	0.8		1.4	
Total**	5.0		5.8	
Total males / females	8.6	1.6	10.1	1.8
WHO Region of the Americas	8.2		8.0	

 $<sup>^{\</sup>star}$  Three-year averages of recorded and unrecorded for 2009–2011 and 2015–2017;  $^{\star\star}$  adjusted for tourist consumption.

Total alcohol per capita (15+) consumption, drinkers only (in litres of pure alcohol), 2016

Beer

Wine

Spirits Other All

	Litres
Males (15+)	19.5
Females (15+)	7.0
Both sexes (15+)	15.2

Prevalence of heavy episodic drinking\* (%), 2016

	Population (15+ years)	Drinkers only (15+ years)	Population (15–19 years)	Drinkers only (15–19 years)
Males	26.4	51.0	20.0	57.0
Females	4.8	18.6	3.2	22.1
Both sexes	15.3	39.9	11.8	47.0

<sup>\*</sup> Consumed at least 60 grams or more of pure alcohol on at least one occasion in the past 30 days.

#### Abstainers (%), 2016

	Males	Females	Both sexes
Lifetime abstainers (15+)	14.8	37.3	26.4
Former drinkers* (15+)	33.5	37.1	35.4
Abstainers (15+), past 12 months	48.3	74.4	61.7

 $<sup>^{\</sup>ast}$  Persons who used to drink alcoholic beverages but have not done so in the past 12 months.

### HEALTH CONSEQUENCES: MORTALITY AND MORBIDITY

Age-standardized death rates (ASDR) and alcohol-attributable fractions (AAF),  $2016\,$ 

	ASI	)R*	AAF	(%)	AAD** (Number)
Liver cirrhosis, males / females	13.6	6.4	59.9	34.5	1 724
Road traffic injuries, males / females	47.3	9.6	31.2	16.8	3 028
Cancer, males / females	180.0	147.4	3.8	1.4	1 447

<sup>\*</sup>Per 100 000 population (15+); \*\*alcohol-attributable deaths, both sexes.

Years of life lost (YLL) score\*, 2016 LEAST < 1 2 3 3 5 > MOST

Prevalence of alcohol use disorders and alcohol dependence (%), 2016\*

	Alcohol use disorders**	Alcohol dependence
Males	10.4	4.5
Females	3.8	1.9
Both sexes	7.0	3.2
WHO Region of the Americas	8.2	4.1

 $<sup>^{\</sup>star}$  12-month prevalence estimates (15+); \*\*including alcohol dependence and harmful use of alcohol.

# POLICIES AND INTERVENTIONS

Written national policy (adopted/revised) / National action plan	Yes (2010/2016) / Yes
Excise tax on beer / wine / spirits	Yes / Yes / Yes
National legal minimum age for off-premise sales of alcoholic beverages (beer / wine / spirits)	18 / 18 / 18
National legal minimum age for on-premise sales of alcoholic beverages (beer / wine / spirits)	18 / 18 / 18
Restrictions for on-/off-premise sales of alcoholic beverages (any): Hours, days / places, density Specific events / intoxicated persons / petrol stations	No, No / No, No No / No / No

National maximum legal blood alcohol concentration (BAC) when driving a vehicle (general / young / professional), in %	0.02 / 0.02 / 0.02
Legally binding regulations on alcohol advertising / product placement (any)	Yes / No
Legally binding regulations on alcohol sponsorship / sales promotion (any)	No / No
Legally required health warning labels on alcohol advertisements / containers (any)	Yes / Yes
National government support for community action (any)	Yes
National monitoring system(s) (any)	Yes

 $<sup>\</sup>ensuremath{^{*}}\xspace$  Based on alcohol-attributable years of life lost.